

previous publications of the Society, it is a book of sterling value, and one that will be of the most essential service to the student of British zoology.

MISCELLANEOUS.

Notes upon the Food of Predaceous Beetles.

By F. M. WEBSTER.

PLINY thought it nothing to the credit of the philosophers of his day that while they were disputing about the number of heroes by the name of Hercules, and the site of the sepulchre of Bacchus, they should not have been able to decide whether or not the queen-bee possessed a sting*.

While the problem of the bee-sting has long been decided, and heroes by the name of Hercules have ceased to trouble the minds of men, there are problems of vital importance regarding the habits of the insects which, during the greater portion of the year, we meet daily in abundance, that still remain unsolved.

The most important as well as the most abundant of these insects are the beetles.

While found in almost every conceivable situation, while our naturalists count the species in their cabinets by thousands, it would be difficult to point out a single species the food-habits of which we *fully* understand, when both the larva and imago state are taken under consideration.

True, we have a sort of ritual laid down by entomologists, based upon the fact that certain species have been known to feed upon certain substances; but this can no more be considered as proof that nothing else enters into their natural diet, than does the meat of which we may partake at dinner prove us to be strictly carnivorous, or the bread or fruit, that we are exclusively vegetarians.

An illustration of this double diet of beetles is found in the case of the European *Silpha opaca*, Linn., the larva of which has been known to feed to an injurious extent upon the leaves of the beet and mangel-wurzel †.

But one of the most fortunate in getting the benefit of our ignorance is the family Carabidæ, to utter a word against which is almost considered a sacrilege.

But, true to the adage "murder will out," occasionally a species is found feeding upon vegetation with a voracity that would do credit to a Chrysomelid. Of these in Europe, besides the *Zabrus gibbus* in both stages, some species of *Pterostichus*, *Anara*, and *Omophron*, and *Calathus latus*, Westw., are said to injure grain by eating off the young shoots or destroying the seed ‡.

Two species of *Bembidium* (*lampos* and *monticola*) have been destructive to the forests of Upper Austria §.

* Plin. Hist. Nat. l. xi. c. 17.

† Curtis, 'Farm Insects,' p. 388.

‡ Report U.S. Agr. Dep. 1868, pp. 79, 80.

§ Deutsche entomologische Zeitschrift, 1879, p. 17.

Brosicus cephalotes attacks the growing grain; and *Aristus bucephalus* devours the seeds of grass*.

In our own country the *Omophron labiatum*, Fab., injures the shoots of young corn in the Southern States.

Harpalus caliginosus, Fab., is suspected of feeding upon grain in stack in Maryland, and also of eating timothy seeds from the heads†.

E. T. Dale, of Jasper, Mo., forwarded to the editors of the 'American Entomologist' specimens of an insect found by him feeding upon the seeds of a plant unknown to him. Upon examination they proved to be *H. caliginosus*‡.

According to Mr. Mather, of Marshalltown, Iowa, the larvæ of some species of *Harpalus* are destructive to his evergreens, he having found them eating off the roots§.

The foregoing is a synopsis of all facts relating to the vegetable-feeding Carabidæ, so far as known to the author of this paper. A number of years ago the writer commenced the study of the food of beetles, correctly judging, from what was then known, that either naturalists were in error in their suppositions, or else that innocent insects were wrongly accused. And he is free to confess his partiality to the former theory as being the most correct. But after several years of study and observation, I have found to my astonishment not only the species accused but others also of this family feeding largely upon vegetable substances, both useful and noxious. Among my earliest observations upon this subject I noted the abundance of Carabidæ about the shocks of wheat in a field where a violent wind-storm had blown down a large number of sheaves, under which, upon their being replaced, large numbers of *Harpalus caliginosus*, *pennsylvanicus*, and *herbivagus*, *Pterostichus lucublandus*, and *Anisodactylus baltimorensis* were observed.

The wheat was drawn in and threshed directly from the field; and a large percentage of the kernels were badly eaten. Previous to the threshing, in another field, a specimen of *H. pennsylvanicus* was captured with a partially eaten grain of wheat in its mandibles. The eaten grains of the threshed wheat seemed to agree with the fragments found in the jaws of the beetle; and as no other destructive elements were noted, the facts seemed to suggest that the damage was done by the before-mentioned Carabidæ. A few days after, *H. pennsylvanicus* was found eating the now fully ripe seeds from a head of upright timothy grass, and was observed to detach them from the glumes. The same species has since been seen feeding largely upon rag-weed (*Ambrosia artemisiifolia*, Linn.) during September, the seeds apparently being the favourite part. A short time after it was found upon timothy grass, it was observed eating the seeds of prairie-grass (*Panicum crus-galli*, L.); and the same day another individual was found devouring an *Ips fasciatus*, Oliv., one of the Nitidulidæ, thus proving its carnivorous propensities also.

* Westwood's Introduction, i. p. 61.

† Report U.S. Agr. Dep. 1868, p. 80.

‡ Am. Ent. o. s. vol. i. p. 80.

§ Am. Ent. n. s. vol. i. p. 26.

H. caliginosus is likewise found eating the seeds of *Ambrosia artemisiæfolia*.

H. herbivagus feeds largely upon the tender shoots of grass during March, cutting them off just below the surface; but later it selects the tender blades and the discoloured parts usually found under boards, &c.

Amara angustata, Say, is found quite abundantly upon the heads of June grass (*Poa pratensis*, L.). But the most voracious Carabid enemy of this grass is the *Anisodactylus sericeus*, Harris.

Early in June 1878 vast numbers of these beetles were noted upon the heads of this grass; in fact, spots several yards in area were literally covered with them. After patient watching (for they are very timid) the proof was conclusive that the unripe seeds were what they were after, and not microscopic insects, as was at first supposed.

The insect is not only cunning, drawing up its legs and dropping to the ground upon the least disturbance, after the manner of a Chrysomelid, but also shows considerable ingenuity. It grasps the lower extremity of the glume tightly in its mandibles, then relaxing slightly, passes upward and again tightens its grasp—a series of movements which finally force the seed, which is now of the consistency of cream, out at the apex. This it at once proceeds to devour with an appetite which reminds one quite forcibly of a tramp who has been obliged to earn his dinner in advance. Later in the season it is found feeding in the same manner upon the seeds of *Agrostis vulgaris*, Witt. Specimens of *Anisodactylus baltimorensis*, Say, were observed feeding upon the marrow and fatty matter clinging to the tibia of some dead animal, probably that of an ox. Attention is called to this as being in perfect accord with microscopic observations reported by Mr. Forbes upon another specimen found upon grass a few months later.

Calathus gregarius, Say, may be found abundantly upon the heads of timothy grass during the early mornings of the beginning of July. Of the genus *Platynus* only a single observation has been obtained; and this was during the latter part of June of the present year, when two specimens of *P. cupripennis*, Say, were seen harassing a half-grown cricket, which they had already disabled. The carnivorous habits of beetles are often as difficult to discover as their vegetarian. Usually they are not at all in favour of public dinners, and, like beasts or birds of prey, prefer to drag their victims to some secluded nook to devour them; hence if the observer gets any insight into this part of their domestic affairs, he must take them by surprise. In this manner a *Staphylinus cinnamopterus*, Grav., was surprised while in the act of devouring an *Anomoglossus pusillus*, Say, having first, to guard against its escape, eaten off four of its legs.

In another instance, a *Dyschirius globulosus*, Say, was observed to spring upon a small salmon-coloured maggot-like larva, and, after disabling it, to start off to select a proper place to devour it. After the lapse of several minutes it returned to drag its victim under a small clod of dirt and leisurely feast upon it.

After the same manner a *Bradycellus rupestris*, Say, was surprised under a stone while eating a small white thread-like worm.

Another family of beetles whose hitherto almost untarnished reputation it seems to have fallen to my lot to soil is the Coccinellidæ. With the exception of *Epilachna borealis*, Fabr., the larva of which feeds upon the vines of the gourd family*, these insects in our country have been considered strictly carnivorous, although several European species are known to deviate from this rule.

This season, specimens of *Megilla maculata*, Deg., have been taken while feeding upon the pollen of the dandelion (*Taraxacum dens-leonis*); and it is not at all improbable that the pollen of other plants also forms a part of their diet, as they are rather common upon the blossoms of plants and fruits.

No accurate estimation of the value of the Coleoptera could be obtained without including the Telephoridæ. Besides *Chauliognathus pennsylvanicus*, Först., which has been found feeding upon the larvæ of the *Conotrachelus nenuphar*, Hbst. †, and *Telephorus bilineatus*, Say, which is such a powerful auxiliary in checking the ravages of the western locust ‡, *Podabrus tomentosus*, Say, has been observed feeding upon the cotton-wood gall-lice, *Pemphigus populiveneræ*, Fitch, and *P. populicaulis*, Fitch. These beetles sometimes place themselves at the opening of the gall, occasionally as many as four together, and catch the mature lice as they attempt an egress, and sometimes plunge their flat head and thorax into the cavity and draw forth and devour large and small indiscriminately. During the latter part of June and the beginning of July these beetles are very abundant, not only upon trees affected by gall-lice, but upon other plants also.—*Illinois State Lab. of Nat. Hist.*, Nov. 1880.

Giant Squid (Architeuthis) abundant in 1875 at the Grand Banks.

By A. E. VERRILL.

From Capt. J. W. Collins, now of the U.S. Fish Commission, I learn that in October 1875 an unusual number of giant squids were found floating at the surface, on the Grand Banks, and mostly entirely dead and more or less mutilated by birds and fishes. In very few cases they were not quite dead, but entirely disabled. These were seen chiefly between N. lat. 44° and 44° 30', and between W. long. 49° 30' and 49° 50'. He believes that between twenty-five and thirty specimens were secured by the fleet from Gloucester, Mass., and that as many more were probably obtained by the vessels from other places. They were cut up and used as bait for codfish. For this use they are of considerable value to the fishermen. Captain Collins was at that time in command of the schooner 'Howard,' which secured five of these giant squids. These were mostly from 10 to 15 feet long, not including the arms, and averaged about 18 inches in diameter. The arms were almost always mutilated. The portion that was left was usually from 3 to 4 feet long, and, at the base, about as large as a man's thigh.

One specimen, when cut up, was packed into a large hogshead

* Am. Ent. o. s. vol. ii. pp. 12 & 373.

† Am. Ent. o. s. vol. i. pp. 35 & 51.

‡ Report U.S. Ent. Com. vol. i. p. 302.